



College of Veterinary Medicine
Department of Internal Medicine & Preventive



Anthrax

Prof. Dr. Maab .AL-Farwachi



College of Veterinary Medicine
Department of Internal Medicine & Preventive



Synonyms:

Anthrax is also called carbúncio or splenic fever



College of Veterinary Medicine

Department of Internal Medicine & Preventive



Definition:

- Anthrax is a rare infectious disease caused by the bacterium *Bacillus anthracis*.
- Anthrax occurs naturally around the world in wild and domestic hoofed animals, especially cattle, sheep, goats, camels .



College of Veterinary Medicine
Department of Internal Medicine & Preventive



How is the disease transmitted and spread?

- Anthrax does typically not spread from animal to animal nor from person to person.



College of Veterinary Medicine
Department of Internal Medicine & Preventive



The bacteria produce spores on contact with oxygen.

- These spores are extremely resistant and survive for years in soil, or on wool or hair of infected animals.



College of Veterinary Medicine

Department of Internal Medicine & Preventive



- Then, if ingested or inhaled by an animal, or on entering through cuts in the skin, they can germinate and cause disease.



College of Veterinary Medicine
Department of Internal Medicine & Preventive



- In carnivores when the animal feeds on an infected source there may be an intestinal form of the disease with fever and cramps from which animals sometimes recover.



College of Veterinary Medicine
Department of Internal Medicine & Preventive



How is the disease diagnosed

- Blood samples from relatively fresh carcasses will contain large numbers of *B. anthracis*, which can be seen under a microscope, cultured or detected by rapid tests, e.g. polymerase chain reaction (PCR).



College of Veterinary Medicine
Department of Internal Medicine & Preventive



- **Transmission and symptoms in humans:**
- It can also occur in humans when they are exposed to the bacterium, usually through handling animals or animal hides



College of Veterinary Medicine
Department of Internal Medicine & Preventive



- **There are three forms affecting humans**
:
- **Cutaneous (skin), inhalation (lungs) and gastrointestinal (stomach and intestine).**



College of Veterinary Medicine
Department of Internal Medicine & Preventive



1- The cutaneous form or skin form is the commonest form and is transmitted via the skin by touch from the carcass, blood, wool, bones or skin of an animal that died of anthrax.



College of Veterinary Medicine
Department of Internal Medicine & Preventive



- **It causes vesicles in the skin that are only slightly painful and black in color. Without treatment, one out of five infected people die.**



College of Veterinary Medicine
Department of Internal Medicine & Preventive



2- The respiratory form is caused by inhalation when working with leathers and wools in a closed atmosphere.



College of Veterinary Medicine
Department of Internal Medicine & Preventive



3 -The intestinal form can be contracted from eating the meat or drinking the blood of an animal affected by this disease. The symptoms of the disease are violent, expressed by vomiting and bloody diarrhea.



College of Veterinary Medicine
Department of Internal Medicine & Preventive





College of Veterinary Medicine
Department of Internal Medicine & Preventive



- **Prevention**
- Any animal that dies suddenly is suspicious of being infected with anthrax: do **not** do an autopsy on an animal that has died suspiciously or has blood oozing out of its natural openings (it can be dangerous)!!



College of Veterinary Medicine
Department of Internal Medicine & Preventive



- * Avoid letting wild dogs and animals open the carcass.**
- * Destroy the carcass as rapidly as possible (by burning or to burying with caustic lime).**



College of Veterinary Medicine
Department of Internal Medicine & Preventive



- * Do not eat meat or blood from an animal that died of an unknown disease.**
- * Be aware of small skin injuries and take care of personal hygiene.**
- ***



College of Veterinary Medicine

Department of Internal Medicine & Preventive



- *Ventilate and use proper working clothes in places where products of animal origin are handled, especially leathers and wool.**
- * Groups of people, who by their work are at serious risk, can be vaccinated.**
- * Vaccinate cattle and goats annually in regions where anthrax is common.**



College of Veterinary Medicine

Department of Internal Medicine & Preventive

